QH1
301
B615

BIOMETRIKA INDEX*

SUPPLEMENT

Author Index: Vols 41–43 (1954–1956)

ARRANGED BY
MAXINE MERRINGTON

[* The last Index published in 1955 contained Author Index: Vols. 1–40 (1901–1953) and Subject Index: Vols. 1–37 (1901–1950).]

ISSUED BY
THE BIOMETRIKA OFFICE, UNIVERSITY COLLEGE, LONDON
PRINTED AT THE UNIVERSITY PRESS, CAMBRIDGE
1957

Printed in Great Britain at the University Press, Cambridge
(Brooke Crutchley, University Printer)

AUTHOR INDEX

Abdel-Aty, S. H. (1954). Tables of generalized k-statistics, **41**, 253–60.

—— (1954). Approximate formulae for the percentage points and the probability integral of the non-central χ^2 distribution, **41**, 538–40

Aitchison, J. & Brown, J. A. C. (1954). An estimation problem in quantitative assay, **41**, 338–43

Andrews, F. C. & Chernoff, H. (1955). A large-sample bioassay design with random doses and uncertain concentration, **42**, 307–15

Anis, A. A. (1955). The variance of the maximum of partial sums of a finite number of independent normal variates, **42**, 96–101

—— (1956). On the moments of the maximum of partial sums of a finite number of independent normal variates, **43**, 79–84

Anscombe, F. J. (1956). On estimating binomial response relations, **43**, 461–4

Anscombe, F. J. & Page, E. S. (1954). Sequential tests for binomial and exponential populations, **41**, 252–3

Arbous, A. G. & Sichel, H. S. (1954). New techniques for the analysis of absenteeism data, **41**, 77–90

Armsen, P. (1955). Tables for significance tests of 2×2 contingency tables, **42**, 494–511

Aty, S. H. Abdel- (1954). Tables of generalized k-statistics, **41**, 253–60

—— (1954). Approximate formulae for the percentage points and the probability integral of the non-central χ^2 distribution, **41**, 538–40

Bailey, N. T. J. (1956). On estimating the latent and infectious periods of measles. Part I. Families with two susceptibles only, **43**, 15–22; Part II. Families with three or more susceptibles, **43**, 322–31

—— (1956). Significance tests for a variable chance of infection in chain-binomial theory, **43**, 332–6

Barnard, G. A. (1954). Simplified decision functions, **41**, 241–51

Bartholomew, D. J. (1954). Note on the use of Sherman's statistic as a test for randomness, **41**, 556–8

—— (1956). A sequential test of randomness for events occurring in time or space, **43**, 64–78

Bartlett, M. S. (1955). Approximate confidence intervals. Part III. A bias correction, **42**, 201–4

Bartlett, M. S. & Medhi, J. (1955). On the efficiency of procedures for smoothing periodograms from time series with continuous spectra, **42**, 143–50

Barton, D. E. (1956). A class of distributions for which the maximum-likelihood estimator is unbiased and of minimum variance for all sample sizes, **43**, 200–2

—— (1956). The limiting distribution of Kamat's test statistic, **43**, 386–7

Barton, D. E. & David, F. N. (1956). Tests for randomness of points on a line, **43**, 104–12

Bechhofer, R. E. *et al.* (1954). A two-sample multiple decision procedure for ranking means of normal populations with a common unknown variance, **41**, 170–6

Blank, A. A. (1956). Existence and uniqueness of a uniformly most powerful randomized unbiased test for the binomial, **43**, 465–6

Bliss, C. I. *et al.* (1956). A rejection criterion based upon the range, **43**, 418–22

Blom, G. (1954). Transformations of the binomial, negative binomial, Poisson and χ^2 distributions, **41**, 302–16 (correction, **43**, 235)

Bose, R. C. (1956). Paired comparison designs for testing concordance between judges, **43**, 113–21

Box, G. E. P. & Hunter, J. S. (1954). A confidence region for the solution of a set of simultaneous equations with an application to experimental design, **41**, 190–9

Bradley, R. A. Rank analysis of incomplete block designs. Part II (1954). Additional tables for the method of paired comparisons, **41**, 502–37; Part III (1955). Some large-sample results on estimation and power for a method of paired comparisons, **42**, 450–70

Broadbent, S. R. (1954). The quotient of a rectangular or triangular and a general variate, **41**, 330–7

—— (1955). Quantum hypotheses, **42**, 45–57

—— (1956). Examination of a quantum hypothesis based on a single set of data, **43**, 32–44

—— (1956). Lognormal approximation to products and quotients, **43**, 404–17

Brown, J. A. C. & Aitchison, J. (1954). An estimation problem in quantitative assay, **41**, 338–43

Butcher, J. C. (1956). Treatment variances for experimental designs with serially correlated observations, **43**, 208–12

Cadwell, J. H. (1954). The statistical treatment of mean deviation, **41**, 12–18

Chanda, K. C. (1954). A note on the consistency and maxima of the roots of likelihood equations, **41**, 56–61

Chapman, D. G. (1955). Population estimation based on change of composition caused by a selective removal, **42**, 279–90

Chernoff, H. & Andrews, F. C. (1955). A large-sample bioassay design with random doses and uncertain concentration, **42**, 307–15

Chu, J. T. (1955). On bounds for the normal integral, **42**, 263–5

—— (1955). The 'Inefficiency' of the sample median for many familiar symmetric distributions, **42**, 520–1

Cochran, W. G. *et al.* (1956). A rejection criterion based upon the range, **43**, 418–22

Cohen, A. C. (1955). Censored samples from truncated normal distributions, **42**, 516–19

Cox, D. R. (1954). The design of an experiment in which certain treatment arrangements are inadmissible, **41**, 287–95

—— (1954). The mean and coefficient of variation of range in small samples from non-normal populations, **41**, 469–81 (correction, **42**, 277)

—— (1956). A note on the theory of quick tests, **43**, 478–80

Cox, D. R. & Smith, W. L. (1954). On the superposition of renewal processes, **41**, 91–9

Cox, D. R. & Stuart, A. (1955). Some quick sign tests for trend in location and dispersion, **42**, 80–95
Crow, E. L. (1956). Confidence intervals for a proportion, **43**, 423–35

Daniels, H. E. (1956). The approximate distribution of serial correlation coefficients, **43**, 169–85
Darwin, J. H. (1956). The behaviour of an estimator for a simple birth and death process, **43**, 23–31
David, F. N. (1955). Studies in the history of probability and statistics. I. Dicing and gaming (a note on the history of probability), **42**, 1–15
—— (1956). A note on Wilcoxon's and allied tests, **43**, 485–8
David, F. N. & Barton, D. E. (1956). Tests for randomness of points on a line, **43**, 104–12
David, F. N. & Johnson, N. L. (1954). Statistical treatment of censored data. Part I. Fundamental formulae, **41**, 228–40
David, F. N. & Kendall, M. G. (1955). Tables of symmetric functions. Part V, **42**, 223–42
David, H. A. (1954). The distribution of range in certain non-normal populations, **41**, 463–8 (correction, **42**, 277)
—— (1955). A note on moving ranges, **42**, 512–15
—— (1956). On the application to statistics of an elementary theorem in probability, **43**, 85–91
—— (1956). Revised upper percentage points of the extreme studentized deviate from the sample mean, **43**, 449–51
David, H. A. *et al.* (1954). The distribution of the ratio, in a single normal sample, of range to standard deviation, **41**, 482–93
Dawson, R. B. (1954). A simplified expression for the variance of the χ^2-function on a contingency table, **41**, 280
Derman, C. (1956). Some asymptotic distribution theory for Markov chains with a denumerable number of states, **43**, 285–94
Douglas, J. B. (1956). Tables of Poisson power moments, **43**, 489
Dunnett, C. W. & Sobel, M. (1954). A bivariate generalization of Student's t-distribution, with tables for certain special cases, **41**, 153–69
—— —— (1955). Approximations to the probability integral and certain percentage points of a multivariate analogue of Student's t-distribution, **42**, 258–60
Dunnett, C. W. *et al.* (1954). A two-sample multiple decision procedure for ranking means of normal populations with a common unknown variance, **41**, 170–6

Fieller, E. C. & Hartley, H. O. (1954). Sampling with control variables, **41**, 494–501
Foster, F. G. (1955). A note on Bailey's and Whittle's treatment of a general stochastic epidemic, **42**, 123–5
Fucks, W. (1954). On Nahordnung and Fernordnung in samples of literary texts, **41**, 116–32

Gani, J. (1955). Some problems in the theory of provisioning and of dams, **42**, 179–200
—— (1955). Some theorems and sufficiency conditions for the maximum-likelihood estimator of an unknown parameter in a simple Markov chain, **42**, 342–59 (correction, **43**, 497–8)
—— (1956). Sufficiency conditions in regular Markov chains and certain random walks, **43**, 276–84
Ghosh, M. N. (1955). Simultaneous tests of linear hypotheses, **42**, 441–9
Ghurye, S. G. & Robbins, H. (1954). Two-stage procedures for estimating the difference between means, **41**, 146–52
Gilbert, N. E. G. (1956). Likelihood function for capture-recapture samples, **43**, 488–9
Good, I. J. (1955). The likelihood ratio test for Markoff chains, **42**, 531–3
Good, I. J. & Toulmin, G. H. (1956). The number of new species, and the increase in population coverage, when a sample is increased, **43**, 45–63
Guest, P. G. (1954). Grouping methods in the fitting of polynomials to equally spaced observations, **41**, 62–76
—— (1956). Grouping methods in the fitting of polynomials to unequally spaced observations, **43**, 149–60
Gulland, J. A. (1955). On the estimation of population parameters from marked members, **42**, 269–70

Haberman, S. (1955). Distributions of Kendall's tau based on partially ordered systems, **42**, 417–24
Haldane, J. B. S. (1955). Substitutes for χ^2, **42**, 265–6
—— (1955). A problem in the significance of small numbers, **42**, 266–7
—— (1955). The rapid calculation of χ^2 as a test of homogeneity from a $2 \times n$ table, **42**, 519–20
Haldane, J. B. S. & Maynard Smith, S. (1956). The sampling distribution of a maximum-likelihood estimate, **43**, 96–103
Hammersley, J. M. & Morton, K. W. (1954). The estimation of location and scale parameters from grouped data, **41**, 296–301
Hannan, E. J. (1955). Exact tests for serial correlation, **42**, 133–42
—— (1955). An exact test for correlation between time series, **42**, 316–26
Hannan, E. J. & Watson, G. S. (1956). Serial correlation in regression analysis. Part II, **43**, 436–48
Harley, B. I. (1954). A note on the probability integral of the correlation coefficient, **41**, 278–80
—— (1956). Some properties of an angular transformation for the correlation coefficient, **43**, 219–24
Hartley, H. O. & Fieller, E. C. (1954). Sampling with control variables, **41**, 494–501
Hartley, H. O. *et al.* (1954). The distribution of the ratio, in a single normal sample, of range to standard deviation, **41**, 482–93
Haskey, H. W. (1954). A general expression for the mean in a simple stochastic epidemic, **41**, 272–5
Healy, M. J. R. (1956). Weighted probits allowing for a non-zero response in the controls, **43**, 207–8
Heine, V. (1955). Models for two-dimensional stationary stochastic processes, **42**, 170–8
Hill, I. D. (1954). The distribution of the regression coefficient in samples from a non-normal population, **41**, 548–52
Hodges, J. L. (1955). Galton's rank-order test, **42**, 261–2
Hoel, P. G. (1954). A test for Markoff chains, **41**, 430–3
Huitson, A. (1955). A method of assigning confidence limits to linear combinations of variances, **42**, 471–9
Hunter, J. S. & Box, G. E. P. (1954). A confidence region for the solution of a set of simultaneous equations with an application to experimental design, **41**, 190–9

Huttly, N. A. (1956). The fitting of regression curves with autocorrelated data, **43**, 468–74
Huzurbazar, V. S. (1955). Exact forms of some invariants for distributions admitting sufficient statistics, **42**, 533–7

Irwin, J. O. (1954). A distribution arising in the study of infectious diseases, **41**, 266–8
Iyer, P. V. Krishna & Kapur, M. N. (1954). Probability distributions arising from points on a line, **41**, 553–4

James, G. S. (1954). Tests of linear hypotheses in univariate and multivariate analysis when the ratios of the population variances are unknown, **41**, 19–43
—— (1955). Cumulants of a transformed variate, **42**, 529–31
—— (1956). On the accuracy of weighted means and ratios, **43**, 304–21
James, G. S. *et al.* (1956). Further critical values for the two-means problem, **43**, 203–5
Jenkins, G. M. (1954). An angular transformation for the serial correlation coefficient, **41**, 261–5
—— Tests of hypotheses in the linear autoregressive model. Part I (1954). Null hypothesis distributions in the Yule scheme, **41**, 405–19; Part II (1956). Null distributions for higher order schemes: non-null distributions, **43**, 186–99
Johnson, N. L. & David, F. N. (1954). Statistical treatment of censored data. Part I. Fundamental formulae, **41**, 228–40
Jonckheere, A. R. (1954). A distribution-free k-sample test against ordered alternatives, **41**, 133–45
Jowett, G. H. (1955). Sampling properties of local statistics in stationary stochastic series, **42**, 160–9

Kamat, A. R. (1954). Distribution theory of two estimates for standard deviation based on second variate differences, **41**, 1–11
—— (1954). Moments of the mean deviation, **41**, 541–2
—— (1956). A two-sample distribution-free test, **43**, 377–85
Kapur, M. N. & Krishna Iyer, P. V. (1954). Probability distributions arising from points on a line, **41**, 553–4
Kendall, M. G. (1954). Note on bias in the estimation of autocorrelation, **41**, 403–4
—— (1954). Two problems in sets of measurements, **41**, 560–4
—— (1956). Studies in the history of probability and statistics. II. The beginnings of a probability calculus, **43**, 1–14
Kendall, M. G. & David, F. N. (1955). Tables of symmetric functions. Part V, **42**, 223–42
Krishna Iyer, P. V. & Kapur, M. N. (1954). Probability distributions arising from points on a line, **41**, 553–4

Lawley, D. N. (1956). Tests of significance for the latent roots of covariance and correlation matrices, **43**, 128–36
—— (1956). A general method for approximating to the distribution of likelihood ratio criteria, **43**, 295–303
Leslie, P. H. (1955). A simple method of calculating the exact probability in 2×2 contingency tables with small marginal totals, **42**, 522–3
Lieblein, J. (1954). Two early papers on the relation between extreme values and tensile strength, **41**, 559–60
Lord, R. D. (1954). The use of the Hankel transform in statistics. Part I. General theory and examples, **41**, 44–55; Part II. Methods of computation, **41**, 344–50

McFadden, J. A. (1956). An approximation for the symmetric, quadrivariate normal integral, **43**, 206–7
Mallows, C. L. (1956). Note on the moment-problem for unimodal distributions when one or both terminals are known, **43**, 224–7
Marriott, F. H. C. & Pope, J. A. (1954). Bias in the estimation of autocorrelations, **41**, 390–402
Mautner, A. J. & Rushton, S. (1955). The deterministic model of a simple epidemic for more than one community, **42**, 126–32
Maynard Smith, S. & Haldane, J. B. S. (1956). The sampling distribution of a maximum-likelihood estimate, **43**, 96–103
Mayne, A. J. (1954). Some further results in the theory of pedestrians and road traffic, **41**, 375–89
Medhi, J. (1956). A note on the risks of error involved in the sequential ratio test, **43**, 231–4
Medhi, J. & Bartlett, M. S. (1955). On the efficiency of procedures for smoothing periodograms from time series with continuous spectra, **42**, 143–50
Mitra, S. K. & Roy, S. N. (1956). An introduction to some non-parametric generalizations of analysis of variance and multivariate analysis, **43**, 361–76
Mohan, C. (1955). The gambler's ruin problem with correlation, **42**, 486–93
Moore, P. G. (1956). The estimation of the mean of a censored normal distribution by ordered variables, **43**, 482–5
Morton, K. W. & Hammersley, J. M. (1954). The estimation of location and scale parameters from grouped data, **41**, 296–301

Nelder, J. A. (1954). The interpretation of negative components of variance, **41**, 544–8

Page, E. S. (1954). Continuous inspection schemes, **41**, 100–15
—— (1955). Control charts with warning lines, **42**, 243–57
—— (1955). A test for a change in a parameter occurring at an unknown point, **42**, 523–7
Page, E. S. & Anscombe, F. J. (1954). Sequential tests for binomial and exponential populations, **41**, 252–3
Patankar, V. N. (1954). The goodness of fit of frequency distributions obtained from stochastic processes, **41**, 450–62
Pearson, E. S. *et al.* (1954). The distribution of the ratio, in a single normal sample, of range to standard deviation, **41**, 482–93
Pillai, K. C. S. (1956). On the distribution of the largest or the smallest root of a matrix in multivariate analysis, **43**, 122–7
Plackett, R. L. (1954). A reduction formula for normal multivariate integrals, **41**, 351–60
Pope, J. A. & Marriott, F. H. C. (1954). Bias in the estimation of autocorrelations, **41**, 390–402
Powell, E. O. (1955). Some features of the generation times of individual bacteria, **42**, 16–44

Quenouille, M. H. (1956). Notes on bias in estimation, **43**, 353–60

Ray, W. D. (1956). Sequential analysis applied to certain experimental designs in the analysis of variance, **43**, 388–403
Reiersøl, O. (1956). A note on the signs of gross correlation coefficients and partial correlation coefficients, **43**, 480–2
Robbins, H. & Ghurye, S. G. (1954). Two-stage procedures for estimating the difference between means, **41**, 146–52
Roy, S. N. & Mitra, S. K. (1956). An introduction to some non-parametric generalizations of analysis of variance and multivariate analysis, **43**, 361–76
Roy, S. N. & Sarhan, A. E. (1956). On inverting a class of patterned matrices, **43**, 227–31.
Royston, E. (1956). Studies in the history of probability and statistics. III. A note on the history of the graphical presentation of data, **43**, 241–7
Ruben, H. (1954). On the moments of order statistics in samples from normal populations, **41**, 200–27 (correction, **41**)
—— (1956). On the sum of squares of normal scores, **43**, 456–8
—— (1956). On the moments of the range and product moments of extreme order statistics in normal samples, **43**, 458–60
Rushton, S. & Mautner, A. J. (1955). The deterministic model of a simple epidemic for more than one community, **42**, 126–32

Sampford, M. R. (1955). The truncated negative binomial distribution, **42**, 58–69
Sarhan, A. E. & Roy, S. N. (1956). On inverting a class of patterned matrices, **43**, 227–31
Shenton, L. R. (1954). Inequalities for the normal integral including a new continued fraction, **41**, 177–89
Sichel, H. S. & Arbous, A. G. (1954). New techniques for the analysis of absenteeism data, **41**, 77–90
Simon, H. A. (1955). On a class of skew distribution functions, **42**, 425–40
Smith, S. Maynard & Haldane, J. B. S. (1956). The sampling distribution of a maximum likelihood estimate, **43**, 96–103
Smith, W. L. & Cox, D. R. (1954). On the superposition of renewal processes, **41**, 91–9
Sobel, M. & Dunnett, C. W. (1954). A bivariate generalization of Student's *t*-distribution, with tables for certain special cases, **41**, 153–69
—— —— (1955). Approximations to the probability integral and certain percentage points of a multivariate analogue of Student's *t*-distribution, **42**, 258–60
Sobel, M. *et al.* (1954). A two-sample multiple decision procedure for ranking means of normal populations with a common unknown variance, **41**, 170–6
Somerville, P. N. (1954). Some problems of optimum sampling, **41**, 420–9
Sterne, T. E. (1954). Some remarks on confidence or fiducial limits, **41**, 275–8
Stuart, A. (1955). A test for homogeneity of the marginal distributions in a two-way classification, **42**, 412–16
Stuart, A. (1955). A paradox in statistical estimation, **42**, 527–9
—— (1956). Bounds for the variance of Kendall's rank correlation statistic, **43**, 474–7
Stuart, A. & Cox, D. R. (1955). Some quick sign tests for trend in location and dispersion, **42**, 80–95
Sundrum, R. M. (1954). On the relation between estimating efficiency and the power of tests, **41**, 542–4

Tate, R. F. (1955). The theory of correlation between two continuous variables when one is dichotomized, **42**, 205–16
Taylor, J. (1956). Exact linear sequential tests for the mean of a normal distribution, **43**, 452–5.
Thompson, H. R. (1954). A note on contagious distributions, **41**, 268–71
—— (1955). Spatial point processes, with applications to ecology, **42**, 102–15
Thomson, G. W. (1955). Bounds for the ratio of range to standard deviation, **42**, 268–9
Toulmin, G. H. & Good, I. J. (1956). The number of new species, and the increase in population coverage, when a sample is increased, **43**, 45–63
Trickett, W. H. & Welch, B. L. (1954). On the comparison of two means: further discussion of iterative methods for calculating tables, **41**, 361–74
Trickett, W. H. *et al.* (1956). Further critical values for the two-means problem, **43**, 203–5
Tukey, J. W. (1955). Interpolations and approximations related to the normal range, **42**, 480–5
Tukey, J. W. *et al.* (1956). A rejection criterion based upon the range, **43**, 418–22

Walker, A. M. (1956). A goodness of fit test for spectral distribution functions of stationary time series with normal residuals, **43**, 257–75
Watson, G. S. (1955). Serial correlation in regression analysis. Part I, **42**, 327–41
—— (1956). On the joint distribution of the circular serial correlation coefficients, **43**, 161–8
—— (1956). A note on the circular multivariate distribution, **43**, 467
Watson, G. S. & Hannan, E. J. (1956). Serial correlation in regression analysis. Part II, **43**, 436–48
Watson, G. S. & Williams, E. J. (1956). On the construction of significance tests on the circle and the sphere, **43**, 344–52
Waugh, W. A. O'N. (1955). An age-dependent birth and death process, **42**, 291–306
Welch, B. L. & Trickett, W. H. (1954). On the comparison of two means: further discussion of iterative methods for calculating tables, **41**, 361–74
Welch, B. L. *et al.* (1956). Further critical values for the two-means problem, **43**, 203–5
Whittle, P. (1954). On stationary processes in the plane, **41**, 434–49 (correction, **42**, 277)
—— (1955). The outcome of a stochastic epidemic—a note on Bailey's paper, **42**, 116–22
—— (1956). On the variation of yield variance with plot size, **43**, 337–43
Wilk, M. B. (1955). The randomization analysis of a generalized randomized block design, **42**, 70–9 (correction, **43**, 235)
Williams, C. B. (1956). Studies in the history of probability and statistics. IV. A note on an early statistical study of literary style, **43**, 248–56

Williams, E. J. (1955). Significance tests for discriminant functions and linear functional relationships, **42**, 360–81

Williams, E. J. & Watson, G. S. (1956). On the construction of significance tests on the circle and the sphere, **43**, 344–52

Williams, R. M. (1956). The variance of the mean of systematic samples, **43**, 137–48

Wise, J. (1955). The autocorrelation function and the spectral density function, **42**, 151–9

—— (1956). Stationarity conditions for stochastic processes of the autoregressive and moving-average type, **43**, 215–19

Wise, M. E. (1954). A quickly convergent expansion for cumulative hypergeometric probabilities, direct and inverse, **41**, 317–29 (correction, **42**, 277)

Wishart, J. (1954). The factorial moments of the distribution of joins between line segments, **41**, 555–6

—— (1956). χ^2 probabilities for large numbers of degrees of freedom, **43**, 92–5

Wooding, R. A. (1956). The multivariate distribution of complex normal variables, **43**, 212–15

Yates, F. (1955). The use of transformations and maximum likelihood in the analysis of quantal experiments involving two treatments, **42**, 382–403

—— (1955). A note on the application of the combination of probabilities test to a set of 2×2 tables, **42**, 404–11